

Stochastic Wobble of Accretion Discs and Jets from Turbulent Rocket Torques

Ryan Pettibone^{1,2} and Eric G. Blackman¹

1. *Department of Physics and Astronomy, University of Rochester, Rochester, NY 14627;*

2. *Present Address: Department of Physics, Caltech, Pasadena, CA, USA, 91125*

ABSTRACT

Models of accretion discs and their associated outflows often incorporate assumptions of axisymmetry and symmetry across the disc plane. However, for turbulent discs these symmetries only apply to averaged quantities and do not apply locally. The local asymmetries can induce local imbalances in outflow power across the disc mid-plane, which can in turn induce local tilting torques. Here we calculate the effect of the resulting stochastic torques on disc annuli that are a consequence of standard mean field accretion disc models. The torques induce a random walk of the vector perpendicular to the plane of each averaged annulus. This random walk is characterized by a radially dependent diffusion coefficient which we calculate for small angle tilt. We use the coefficient to calculate a radially dependent time scale for annular tilt and associated jet wobble. The wobble time depends on the square of the wander angle so the age of a given system determines the maximum wobble angle. We apply this to examples of blazars, young stellar objects and binary engines of pre-planetary nebulae and microquasars. It is noteworthy that for an averaging time $t_w \sim 3$ days, we estimate a wobble angle for jets in SS433 of $\theta \sim 0.8$ degrees, not inconsistent with observational data. In general the non-periodic nature of the stochastic wobble could distinguish it from faster periodic jet precession.

1. Introduction

Winds and jets are commonly associated with accretion disc engines, and standard models of discs (Shakura & Sunyaev 1973; Pringle 1981; Frank et al. 2002) and outflows (Blandford & Payne 1982; Pelletier & Pudritz 1992) typically impose axisymmetry and reflection symmetry across the disc plane. However, theory and observation suggest that for a wide range of sources, angular momentum transport is likely mediated at least in part by turbulence (e.g. Terquem (2002)). This immediately implies that the assumptions of

axisymmetry apply at most only to mean flow quantities, averaged over sufficient time or space (e.g. Balbus et al. 1994). The local violation of symmetries for a turbulent disc, leads to an intrinsic variability (Blackman 1998).

We suggest that an additional consequence of the fluctuations may be a local “rocket effect.” When a wind is expelled from the surface of a disc it exerts a force on that surface which compensates for the momentum of the outflow. In idealized models of symmetric accretion discs the forces exerted on the disc by the winds on either side of the disc cancel out. In reality, although these forces must cancel globally if the disc center of mass is stationary, turbulence induced asymmetry makes it unlikely that these forces cancel out locally. Thus, different portions of the disc can be displaced above or below the initial mean symmetry plane. Here we investigate how disc turbulence can cause a local imbalance between the winds emanating from the top and bottom by averaging the local effect over azimuth. The net effect is to induce a stochastic wander of the vector perpendicular to the plane determined by the azimuthal average of the asymmetry for each eddy at a given radius. We model this as Brownian motion, and calculate the associated diffusion coefficient.

In Sec. 2 we discuss the needed basics of Brownian motion. In Sec. 3 we then derive the equation for the aforementioned stochastic diffusion coefficient for tilt as a function of disc parameters. In Sec. 4 we apply this diffusion coefficient to the wobbling of a thin accretion disc and a stochastic wander of the associated jet axis, assumed to be parallel to the disc’s primary angular velocity vector. We apply the formalism to four astrophysical engine classes (blazars, young stellar objects (YSOs), planetary nebulae, microquasars) and estimate the wander angles for which the wobble could be observable in these systems. As a particular example, we compare the crudely predicted wobble to that observed in SS433 and find agreement to order of magnitude for a plausible choice of disc parameters. The stochastic nature of the tilt distinguishes it from that of an ordered precession. We conclude in Sec. 5.

2. Review of Brownian Motion

Consider the random walk on a two-dimensional square lattice of points (Chandrasekhar 1943). We label a point on the lattice by coordinates (θ, ϕ) , and the spacing between adjacent lattice points by $\Delta\theta$. We also break time into discrete intervals Δt . We use these coordinates in anticipation of our application to the random walk of a unit vector tip on the surface of a sphere, locally approximated as a plane for small angles.

At time $t - \Delta t$, a point particle located at (θ, ϕ) moves by time t to one of the 4 points

$(\theta + \Delta\theta, \phi)$, $(\theta - \Delta\theta, \phi)$, $(\theta, \phi + \Delta\theta)$, $(\theta, \phi - \Delta\theta)$ with equal probability. If the particle moves *into* the lattice point (θ, ϕ) at time t , it would have come from one of the four adjacent lattice points in the previous time step. Therefore, if $P(\theta, \phi, t)$ is the probability the particle is located at (θ, ϕ) at time t , we have

$$P(\theta, \phi, t) = \frac{1}{4}(P(\theta + \Delta\theta, \phi, t - \Delta t) + P(\theta - \Delta\theta, \phi, t - \Delta t) + P(\theta, \phi + \Delta\theta, t - \Delta t) + P(\theta, \phi - \Delta\theta, t - \Delta t)) \quad (1)$$

This allows us to write a difference equation for $P(\theta, \phi, t)$. Using (1) we have

$$\frac{P(\theta, \phi, t) - P(\theta, \phi, t - \Delta t)}{\Delta t} = D(A(\theta, \phi) + B(\theta, \phi)) \quad (2)$$

where

$$A(\theta, \phi) = \frac{P(\theta + \Delta\theta, \phi, t - \Delta t) - 2P(\theta, \phi, t - \Delta t) + P(\theta - \Delta\theta, \phi, t - \Delta t)}{(\Delta\theta)^2}, \quad (3)$$

$$B(\theta, \phi) = \frac{P(\theta, \phi + \Delta\theta, t - \Delta t) - 2P(\theta, \phi, t - \Delta t) + P(\theta, \phi - \Delta\theta, t - \Delta t)}{(\Delta\theta)^2}, \quad (4)$$

and

$$D = \frac{(\Delta\theta)^2}{4\Delta t}. \quad (5)$$

Equation (2) is a discrete version of the two-dimensional diffusion equation that characterizes a random walk with diffusion coefficient D . Although we will use the discrete version in this paper, a continuum diffusion equation results by taking the continuum limit for which $\Delta\theta \rightarrow 0$ and $\Delta t \rightarrow 0$. In this limit, $A \rightarrow \frac{\partial^2 P}{\partial \theta^2}$ and $B \rightarrow \frac{\partial^2 P}{\partial \phi^2}$. If we then define $D \equiv \lim_{\Delta t, \Delta\theta \rightarrow 0} (\Delta\theta)^2 / \Delta t$, we have:

$$\frac{\partial P}{\partial t} = D \left(\frac{\partial^2 P}{\partial \theta^2} + \frac{\partial^2 P}{\partial \phi^2} \right) \quad (6)$$

which is a two-dimensional Cartesian diffusion equation in the local angle coordinates.

3. Derivation of the Wobble Diffusion Coefficient

To derive D for a disc, we first consider a small volume element at radius r with azimuthal and radial scales equal to that of a turbulent eddy and with vertical scale equal to the disc thickness. We define dL as the net mechanical wind luminosity emanating from this

volume element, and v as the associated mean outflow speed at the chosen radius. We posit that the winds emanating from either side of thin annulus are not exactly in momentum balance at all times due to turbulent fluctuations. We denote wind mechanical luminosities for a given eddy emanating from the top and bottom of the disc as dL_{top} and dL_{bot} respectively, such that $dL = dL_{top} + dL_{bot}$. Let ϵ_{top} and ϵ_{bot} denote unit vectors in the direction of wind flow on the top and bottom. The net force on the volume element from the two winds is

$$d\mathbf{F} = d\mathbf{F}_{top} + d\mathbf{F}_{bot} = \frac{dL_{top}}{v}\epsilon_{top} + \frac{dL_{bot}}{v}\epsilon_{bot} = \epsilon' \frac{dL}{v}, \quad (7)$$

where $\epsilon' \equiv (\epsilon_{top}dL_{top} + \epsilon_{bot}dL_{bot})/dL$, and characterizes the direction of wind imbalance. Since we have assumed that the origin of the force imbalance (turbulence) is intrinsically random, ϵ' will be a random vector dependent on both time and position. Fluctuations around both the mean outflow speed and mass outflow rate can be incorporated into ϵ' given its definition; the explicitly presence of the mean velocity in (7) does not preclude this.

The force on a disc volume element results from a “rocket effect” created by non-vanishing last term in (7). To calculate the associated torque, we assume that ϵ' is a poloidal vector ($\epsilon' \cdot \hat{\phi} = 0$) for every volume element. The associated torque then lies in the initial disc plane and is given by

$$d\boldsymbol{\tau} = (\mathbf{r} \times d\mathbf{F}) = (\mathbf{r} \times \epsilon') \frac{dL}{v} = (\hat{\mathbf{r}} \times \epsilon') \frac{r}{v} dL. \quad (8)$$

Summing the torques for all volume elements at radius r leaves a net torque on an annulus of radial thickness dr at radius r given by

$$d\boldsymbol{\tau}_{net} = \sum_{dV \in E(r)} (\hat{\mathbf{r}} \times \epsilon') \frac{r}{v} dL = \epsilon \frac{r}{v} \frac{dL}{dr} dr, \quad (9)$$

where we have defined $E(r) \equiv \{\text{all volume elements at radius } r\}$, and $\epsilon \equiv \sum (\hat{\mathbf{r}} \times \epsilon')$. Note that since ϵ' is poloidal, $\hat{\mathbf{r}} \times \epsilon'$ lies in the plane of the disc, and therefore so does ϵ , being a sum of such vectors. Since $d\boldsymbol{\tau}$ lies in the plane of the disc, so does $d\boldsymbol{\tau}_{net}$. By summing over all the volume elements in the annulus we have removed the azimuthal dependence, providing what is needed to characterize the response of an axisymmetric mean field annulus. This is an annulus of a “standard” axisymmetric radially dependent accretion disc (which itself is axisymmetric precisely because of the azimuthal average over turbulent fluctuations).

We can now calculate the rate of change of angular momentum of this annulus associated with tilting from its initial disc symmetry plane. This tilt introduces an angular momentum vector in the initial disc plane that bisects the annulus, and is approximately perpendicular to the underlying Keplerian rotation vector. The component of the moment of inertia tensor

appropriate for the tilt axis is $dI = \frac{1}{2}r^2 dm(r)$, where the mass of the annulus is $dm(r) = \mu(r)dr$ and $\mu(r)$ is the mass per unit radius. The rate of change in angular momentum that characterizes the tilt is then

$$d\dot{\mathbf{J}} = \frac{d\boldsymbol{\omega}}{dt}dI = \frac{r^2}{2} \frac{d\boldsymbol{\omega}}{dt} dm = \frac{r^2}{2} \frac{d\boldsymbol{\omega}}{dt} \mu(r) dr, \quad (10)$$

where $\boldsymbol{\omega}$ is the orbital angular momentum vector that would reflect a fixed-plane orbital motion of disc material in the absence of rocket torques. Using Newton's second law $d\dot{\mathbf{J}} = d\boldsymbol{\tau}_{net}$. It follows that

$$\frac{r^2}{2} \frac{d\boldsymbol{\omega}}{dt} \mu(r) = \boldsymbol{\epsilon} \frac{r}{v} \frac{dL}{dr}. \quad (11)$$

If we assume that $\boldsymbol{\epsilon}$ varies on a characteristic time scale Δt , then (11) implies

$$\Delta\boldsymbol{\omega} = \frac{2}{rv\mu(r)} \frac{dL}{dr} \boldsymbol{\epsilon} \Delta t. \quad (12)$$

Eq. (12) measures the tilt of the plane the annulus. Since $d\boldsymbol{\tau}_{net}$ is in the plane of the annulus, the torque has the effect of rotating $\boldsymbol{\omega}$ but not changing its magnitude. If the angular deflection is small in a time Δt , then $\Delta\theta \sim \frac{|\Delta\boldsymbol{\omega}|}{\omega}$, where $\Delta\theta$ is the change in the angle of $\boldsymbol{\omega}$ during the time Δt . We can then appeal to the Cartesian lattice calculation of Sec. 2 to obtain the small angle diffusion coefficient: Combining the expression for $\Delta\theta$ just derived with (12) and (5) we obtain

$$D(r) = \frac{1}{r^2 \mu(r)^2 v^2 \omega^2} \left(\frac{dL}{dr} \right)^2 \epsilon^2 \Delta t = \frac{1}{r^2 \mu(r)^2 v^2 \omega^2} \left(\frac{dL}{dr} \right)^2 \frac{\epsilon_{ed}^2 t_{ed}}{\sqrt{N}}, \quad (13)$$

where $\epsilon_{ed} = N^{1/2} \epsilon$ represents the characteristic contribution from each of the N eddies at a given radius to ϵ , and $\tau_{ed} = \tau/N^{1/2}$ is the characteristic time scale for evolution of a given eddy. Here $N = 2\pi r/l_{ed}$ is the number of eddies with scale l_{ed} at radius r .

This diffusion coefficient characterizes the random walk of the orbital velocity vector of the mean field annulus due to tilt from the stochastic rocket force. Since the magnitude of $\boldsymbol{\omega}$ never changes in our approximation, the tilt implies that the orbital angular vector moves on the surface of a sphere as the annulus wobbles. For small tilt angles, this surface can be locally approximated by a plane and we are justified in using the Cartesian formulation of Sec. 2, despite the global differences between spheres and planes.

The derivation and use of Eqn. (13) implies that each radius has its own wobble diffusion coefficient. This does not violate angular momentum conservation because annuli are (at least) viscously coupled. If all of the material that enters the disc at large radii is coplanar, then conservation of total angular momentum implies that the total integrated

angular momentum about any axis perpendicular to that of the initial ω must vanish at all times. This means that if some inner mean annulus of gas tilts in one direction, there will be a corresponding counter-tilt elsewhere. Turbulent viscosity allows the radial distribution of tilts to be non-trivial and because jet power is likely dominated by inner annuli, the tilt of the inner-most annuli are particularly relevant for predicting jet wobble because they dominate the accretion power. In what follows, we assume that the axis of any annular section of a jet tracks the orbital axis of the corresponding disc annulus to which that jet section is anchored. Thus the jet wobble directly tracks the disc wobble.

4. Application to Astrophysical Engines

Here we apply the above formulae to thin accretion discs and jet wobble.

4.1. Determination of Stochastic Wobble times for Discs and Jets

For a thin, non-self-gravitating disc in hydrostatic equilibrium

$$\omega = \omega_K \equiv \sqrt{GM/r^3} \sim c_s/H \sim t_{ed}^{-1} \quad (14)$$

where c_s is the sound speed, M is the central object mass, and H is the density scale height. The Shakura-Sunyaev (Shakura & Sunyaev (1973)) viscosity ν in hydrostatic equilibrium then satisfies $\nu \equiv \alpha c_s H \sim v_{ed}^2/t_{ed} \sim v_{ed}^2 H/c_s$, where α is the viscosity parameter. Then (e.g. Blackman 1998) $v_{ed} \sim \alpha^{1/2} c_s$ and

$$l_{ed} = \alpha^{1/2} H, \quad (15)$$

so

$$N = \frac{2\pi}{\alpha^{1/2}} \frac{r}{H}. \quad (16)$$

The disc accretion rate can be modeled as $\dot{M}_a(r) = \dot{M}_o(r/r_o)^s$ (e.g. Blandford & Begelman (1999)), where $\dot{M}_o$ is the accretion rate at the outer edge of the disc, r_o is the radius of the outer edge, and $0 < s < 1$ is a parameter. The total mass outflow rate from r_o to r is then $\dot{M}(r) = \dot{M}_o[1 - (r/r_o)^s]$. Taking the derivative, we obtain the outflow mass loss rate by a small annulus of width dr at radius r to be

$$\frac{d\dot{M}}{dr} dr = -\dot{M}_o \frac{s}{r_o} \left(\frac{r}{r_o}\right)^{s-1} dr. \quad (17)$$

The luminosity of the gas ejected by the wind in the annulus is $dL = \frac{1}{2}v^2|d\dot{M}|$, where the outflow speed satisfies

$$v \simeq v_{esc} \equiv \sqrt{2GM/r}, \quad (18)$$

so in combination with (17) we have

$$\frac{dL}{dr} = \dot{M}_o \frac{s}{2r_o} \left(\frac{r}{r_o}\right)^{s-1} v_{esc}^2. \quad (19)$$

Using (14), (15), (16), (18) and (19) in (13) gives

$$D(r) = 4.4 \times 10^{19} \alpha^{1/4} \left(\frac{s\epsilon_{ed}}{\mu}\right)^2 \left(\frac{H}{r_g}\right)^{\frac{1}{2}} \left(\frac{M}{M_\odot}\right)^{-1} \left(\frac{r}{r_o}\right)^{2s-1} \left(\frac{r_o}{r_g}\right)^{-1} \left(\frac{\dot{M}_o}{10^{18} \text{ g/s}}\right)^2 \text{ rad}^2/\text{s}, \quad (20)$$

where $r_g = GM/c^2$. For a standard Shakura-Sunyaev disc model (Shakura & Sunyaev 1973) supplemented by our radially dependent accretion rate $\dot{M} = \dot{M}_o(r/r_o)^s$ we have

$$\mu(r) = 5.1 \times 10^{11} \alpha^{-4/5} \left(\frac{\dot{M}_o}{10^{18} \text{ g/s}}\right)^{7/10} \left(\frac{r}{r_o}\right)^{7s/10} \left(\frac{M}{M_\odot}\right)^{1/2} \left(\frac{r}{r_g}\right)^{1/4} \left[1 - \left(\frac{r_g}{r}\right)^{1/2}\right]^{7/20} \text{ g/cm}, \quad (21)$$

and

$$\frac{H}{r_g} = 9 \times 10^{-3} \alpha^{-1/10} \left(\frac{M}{M_\odot}\right)^{-1/4} \left(\frac{\dot{M}_o}{10^{18} \text{ g/s}}\right)^{3/20} \left(\frac{r}{r_o}\right)^{3s/20} \left(\frac{r}{r_g}\right)^{9/8} \left[1 - \left(\frac{r_g}{r}\right)^{1/2}\right]^{3/20}. \quad (22)$$

Using (21) and (22) in (20), the time required to wander an angle θ radians is

$$t_w(r) = \frac{\theta^2}{4D} = 1.6 \times 10^4 \alpha^{-9/5} \left(\frac{\theta}{s\epsilon_{ed}}\right)^2 \left(\frac{\dot{M}_o}{10^{18} \text{ g/s}}\right)^{\frac{-27}{40}} \left(\frac{r}{r_g}\right)^{\frac{17}{16} - \frac{27s}{40}} \left(\frac{M}{M_\odot}\right)^{17/8} \left(\frac{r_o}{r_g}\right)^{27s/40} \left[1 - \left(\frac{r_g}{r}\right)^{1/2}\right]^{5/8} \text{ sec}. \quad (23)$$

For stochastic wobble to be observable $t_w(\theta)$ must at least be less than the age of a system τ_{age} . A further constraint follows from the standard accretion paradigm in which accretion supplies the energy for a turbulent viscosity which, together with the outflow, transport angular momentum and sustain the accretion. As discussed at the end of Sec.3, the turbulent viscosity is also a means of transporting the angular momentum associated with tilt between annuli such that the tilt of inner annuli is compensated by the opposing tilt of outer annuli. For stochastic wobble to produce a net mean tilt on an annulus at an inner radius, the viscous time scale there $\tau_\nu(r)$ should be short compared to the tilt time, so that the compensating angular momentum opposite to that associated with the tilt can be transported outward. Note that even though material also accretes on a viscous time, the resupply of material at a given radius would also maintain the stochastic equilibrium on time

scales short compared to the secular wobble time. Provided there is a supply of mass, we therefore consider viscosity to maintain the conditions for the effect to accrue, rather than washing out the effect at a given radius.

Summarizing the above conditions to be checked, we have

$$\tau_v(r) < t_w(r) < \tau_{age}, \quad (24)$$

where, from (22), the viscous accretion time scale for $r > 3r_g$ is given by

$$\tau_v(r) \simeq \frac{r^2}{\alpha H^2 \Omega} = 6 \times 10^{-2} \alpha^{-4/5} \left(\frac{M}{M_\odot} \right)^{3/2} \left(\frac{\dot{M}_0}{10^{18} \text{ g/s}} \right)^{-3/10} \left(\frac{r}{r_g} \right)^{-\frac{3s}{10} + \frac{5}{4}} \left(\frac{r_0}{r_g} \right)^{3s/10} \text{ sec}, \quad (25)$$

and

$$\tau_{age} = \text{Max} \left[\tau_v(r_o), \frac{M_c}{\dot{M}} \right] \quad (26)$$

where M_c is the companion mass. The second value in brackets is used when the accretion disc is being supplied by a companion, while the first (the viscous time at the outer disc radius) is used when there is no companion-fed resupply.

For later use in evaluating (24), we have from (23) and (25)

$$\frac{t_w(r)}{\tau_v(r)} \sim 2.7 \times 10^5 \alpha^{-1} \left(\frac{\theta}{s\epsilon_{ed}} \right)^2 \left(\frac{\dot{M}_0}{10^{18} \text{ g/s}} \right)^{-3/8} \left(\frac{r}{r_g} \right)^{-\frac{3}{16} - \frac{3s}{8}} \left(\frac{M}{M_\odot} \right)^{5/8} \left(\frac{r_0}{r_g} \right)^{3s/8} \quad (27)$$

and

$$\frac{t_w(r)}{\tau_v(r_o)} \sim 2.7 \times 10^5 \alpha^{-1} \left(\frac{\theta}{s\epsilon_{ed}} \right)^2 \left(\frac{\dot{M}_0}{10^{18} \text{ g/s}} \right)^{-3/8} \left(\frac{r}{r_g} \right)^{\frac{17}{16} - \frac{27s}{40}} \left(\frac{M}{M_\odot} \right)^{5/8} \left(\frac{r_0}{r_g} \right)^{\frac{27s}{40} - \frac{5}{4}} \quad (28)$$

Note that the left side of (28) employs r for the numerator and r_o in the denominator on the left hand side as that ratio represents a comparison between the wobble time at a given radius with the accretion time at the outer radius. Eq. (27) compares the two time scales at the same radius. In the next section we estimate (23) and check condition (24) for example classes of accretion engines.

4.2. Application to Astrophysical Jets

To find the characteristic jet/disc wobble times for various systems, we now estimate (23) for active galactic nuclei (AGN) young stellar objects (YSO), microquasars, planetary nebulae (PNe) and active galactic nuclei (AGN). Since the wander time scale depends on the square of the wander angle, we can determine the wander angle for which the observability condition (25) is satisfied. Presently, the quantities s and ϵ_{ed} , and α are not tightly pinned

by either theory or simulation so we maintain their dependence explicitly. Fiducial values will be taken to be $\epsilon_{ed} = 0.5$, and a mass-loss power law index $s = 0.2$, so their product can be scaled to 0.01. We also scale the results with α .

Consider first AGN blazars (e.g. 3C273; Paltani and Türler 2005). We use $M \sim 10^9 M_\odot$ and $\dot{M}_0 \sim 10 M_\odot/\text{yr}$. Then taking $r_0 = 10^6 r_g$ and $r = r_g \sim 10^{14} \text{cm}$, we find that $t_w \sim 3.1 \times 10^{11} \left(\frac{s\epsilon_{ed}}{0.01}\right)^{-2} \left(\frac{\alpha}{0.1}\right)^{-9/5} \left(\frac{\theta}{0.03 \text{ rad}}\right)^2 \text{yr}$. For these parameters the condition $\theta < 0.03$ corresponds to satisfying the second inequality in (24), namely that Eq. (28) is ≤ 1 . However, observable duty cycles of AGN jets seem to last only $\sim 10^8 \text{yr}$, which would requires a wobble angle $\theta < 100''$. For this latter range of wobble angles, both inequalities in (24) are satisfied, using the first value on the right of (26).

For YSOs (Hartmann 1998), we take $M = 1 M_\odot$, $\dot{M} \sim 10^{-5} M_\odot/\text{yr}$, $r_o = 100 \text{AU}$. At $r = 7 \times 10^7 r_g = 10^{13} \text{cm}$, Eq. (23) then gives $t_w \sim 4.6 \left(\frac{s\epsilon_{ed}}{0.01}\right)^{-2} \left(\frac{\alpha}{0.3}\right)^{-9/5} \left(\frac{\theta}{0.03 \text{ rad}}\right)^2 \text{yr}$. Using (27), the first inequality of (24) is easily satisfied. For the second inequality of (24) we use the first value on the right of (26). From (28) we then find $\frac{t_w(r)}{\tau_v(r_o)} \sim 0.3 \left(\frac{\alpha}{0.3}\right)^{-1} \left(\frac{\theta}{0.01}\right)^2$ satisfied for wander angles $\leq 0.01 \text{rad}$.

Accretors with binary companions supply accretion for times much longer than the viscous time scale at the outer radius. This makes the second equality in (24) easier to satisfy for large wobble angles in those systems. Two examples of such are accretors are pre-PNe engines and microquasars. For pre-PNe engines, an accretion disc can form around the post-ascending giant branch (AGB) white dwarf core as the result of common envelope evolution (Nordhaus & Blackman 2006). We take $M = 0.6 M_\odot$, $\dot{M} \sim 10^{-5} M_\odot/\text{yr}$, $r_o = 10^{11} \text{cm}$. Here the engine accretor is a proto-white dwarf of radius $\sim 10^9 \text{cm}$ or $r \sim 10^4 r_g$. At $r = 10^5 r_g \sim 10^{10} \text{cm}$, Eq. (23) then gives $t_w \sim 5.5 \left(\frac{s\epsilon_{ed}}{0.01}\right)^{-2} \left(\frac{\alpha}{0.3}\right)^{-9/5} \left(\frac{\theta}{0.03 \text{ rad}}\right)^2 \text{yr}$. Again for this case the first inequality in (24) is satisfied from (27): At the inner disc radius $t_w/\tau_\nu \sim 700 \left(\frac{\theta}{0.03 \text{ rad}}\right)^2$. In this case however, the second equality in (24) is more easily satisfied as we can use the second value in (26): The pre-PNe outflows last $\tau_{age} \sim 10^3 \text{yr}$ and the same accretion disc, supplied by a very low mass star, or very high mass planet, may in fact be continuously supplying outflow power even during the 10^4yr PNe phase (Blackman et al. 2001). Both inequalities of (24) can then be satisfied for $\theta \leq 0.8 \text{rad}$.

For microquasar examples we consider Cygnus X-1 (Young et al. 2001) and SS433 (Katz & Piran 1982; Begelman et al. 2006; Blundell et al. 2005; 2007). For the former, we take $M = 10 M_\odot$, $\dot{M} \sim 10^{19} M_\odot/\text{yr}$, and $r_o = 1000 r_g$. At $r = r_g \sim 1.5 \times 10^6 \text{cm}$, Eq. (23) then gives $t_w \sim 0.03 \left(\frac{s\epsilon_{ed}}{0.01}\right)^{-2} \left(\frac{\alpha}{0.3}\right)^{-9/5} \left(\frac{\theta}{0.03 \text{ rad}}\right)^2 \text{yr}$. For this case too, the first inequality in (24) is easily satisfied from (27). The second inequality in (24) allows use of the second value in (26). The companion mass to e.g. Cyg X-1 is at least $20 M_\odot$ so for accretion at $10^{-7} M_\odot/\text{yr}$, accretion can be powered long after the wobble time.

For SS433 (e.g. Begelman et al. 2006) $M = 10M_{\odot}$, and we take the inner radius $r \sim r_g \sim 1.5 \times 10^6 \text{cm}$. The accretion rate that makes it to the inner region, relevant for the jets, is $\dot{M}_{in} \sim 2 \times 10^{19} \text{g/s}$. This is to be distinguished from the super-Eddington accretion rate of $\dot{M}_{out} \geq 10^3 \dot{M}_{in}$ at the disc circularization radius of $r_{circ} \sim 10^{12} \text{cm}$. Begelman et al. (2006) argue that the much of the mass is taken out by a wind at $r \sim \frac{r_{circ}}{10} \sim 10^5 r_g \sim 10^{11} \text{cm}$. We therefore take $r_o \sim 10^5 r_g$ for the outer radius of the disc supplying mass to the inner jet and use $\dot{M}_{in}$ for the relevant accretion rate. We then obtain $t_w \sim 0.04 \left(\frac{s\epsilon_{ed}}{0.01}\right)^{-2} \left(\frac{\alpha}{0.3}\right)^{-9/5} \left(\frac{\theta}{0.03 \text{ rad}}\right)^2$ yr. For SS433, like Cygnus X-1, the first inequality in (24) is easily satisfied from (27) and a companion mass of $\sim 20M_{\odot}$ (based on crudely averaging recent mass estimates of Lopez et al. 2006 and Hillwig and Gies 2008) allows use of the second value in (26) for the second inequality in (24). This implies that accretion will be fueled long after the wobble time. Again, we have not considered relativistic effects.

We can compare the stochastic wander just estimated for SS433 with the data of Blundell et al. (2005,2007), who monitored fluctuations in the speed and direction of the inner jets. Table 1 of Blundell et al. (2007) shows that when averaged over several day time scales, wander angles of $\sim 0.5 - 0.7$ degrees were obtained. This is roughly consistent with our result above: Converting units, our result for SS433 can be written $\theta \sim 1.7 \left(\frac{t_w}{15 \text{days}}\right)^{1/2} \text{deg}$.

There is likely a maximum time scale $t_{w,max}$ which is the value of t_w above which the stochastic wander is tempered by some negative feedback from forces not considered in the present work. Thus $t_w \leq t_{w,max}$ if the wander is to be calculated with our approximations. One limit on t_w comes from the small wander angle approximation that allowed our use of a 2-D Cartesian approximation. However, for a precessing and nodding system like SS433 (e.g. Fabrika 2004), the 162 day precession time scale or the 6.3 day nodding time scale may provide the maximum t_w appropriate for computing untempered stochastic wander. We do not determine $t_{w,max}$ here or study large angle wobble but note that a persistent feature of stochastic wander is its non-periodicity, which distinguishes it from systematic precession induced by binary companions (e.g. Terquem et al. 1999).

5. Conclusions

Turbulence in accretion discs violates local axisymmetry and reflection symmetry even if these properties hold in some average mean sense. As such, this symmetry violation likely also applies to outflows emanating from the disc. Due to a rocket-like effect, any outflow power asymmetry locally displaces the disc material in the direction opposite to the net outflow. By azimuthally averaging outflows with locally reflection asymmetric powers, we have calculated the radially dependent small-angle diffusion coefficient of the angular

momentum vector for disc annuli. The diffusion of this vector represents a stochastic tilting of the annulus defined by the averaging, and implies a disc warping and jet wobble. For the latter, the most relevant tilt is that associated inner disc radii. We have calculated estimated the stochastic wobble angle vs. time scale for jets and applied the relation to a handful of astrophysical engine classes to exemplify crude application of the paradigm. The numbers are not inconsistent with measured values in SS433. We have not considered the nonlinear saturation of the stochastic wobble in the present work.

EGB acknowledges support from NSF grant AST-0406799, NASA grant ATP04-0000-0016, and NORDITA during the Turbulence and Dynamos workshop 2008.

REFERENCES

- Balbus, S. A., Gammie, C. F., & Hawley, J. F. 1994, MNRAS, 271, 197
- Begelman, M. C., King, A. R., & Pringle, J. E. 2006, MNRAS, 370, 399
- Blandford, R. D., Begelman, M. C., 1999, MNRAS, 303, L1
- Blandford, R. D., Payne, D. G., 1982, MNRAS, 199, 883
- Blackman, E. G., Frank, A., & Welch, C. 2001, ApJ, 546, 288
- Blackman, E. G. 1998, MNRAS, 299, L48
- Blundell, K. M., & Bowler, M. G. 2005, ApJ, 622, L129
- Blundell, K. M., Bowler, M. G., & Schmidtobreick, L. 2007, A&A, 474, 903
- Chandrasekhar, S., 1943, Rev. Mod. Phys. 15, 1
- Collins, G. W., II, & Newsom, G. H. 1986, ApJ, 308, 144
- Collins, G. W., & Scher, R. W. 2002, MNRAS, 336, 1011
- Eikenberry, S. S., Cameron, P. B., Fierce, B. W., Kull, D. M., Dror, D. H., Houck, J. R., & Margon, B. 2001, ApJ, 561, 1027
- Fabrika, S. 2004, Astrophysics and Space Physics Reviews, 12, 1
- Fender, R. P., Stirling, A. M., Spencer, R. E., Brown, I., Pooley, G. G., Muxlow, T. W. B., & Miller-Jones, J. C. A. 2006, MNRAS, 369, 603

- Frank, J., King, A., & Raine, D. J. 2002, *Accretion Power in Astrophysics*, by Juhan Frank and Andre w King and Derek Raine, pp. 398. ISBN 0521620538. Cambridge, UK: Cambridge Unive rsity Press, February 2002.,
- Hartmann, L., *Accretion Processes in Star Formation*, 1998, (Cambridge Univ. Press: Cambridge UK)
- Hillwig, T. C., & Gies, D. R. 2008, *ApJ*, 676, L37
- Lopez, L. A., Marshall, H. L., Canizares, C. R., Schulz, N. S., & Kane, J. F. 2006, *ApJ*, 650, 338
- Katz, J. I., & Piran, T. 1982, *Astrophys. Lett.*, 23, 11
- Margon, B., & Anderson, S. F. 1989, *ApJ*, 347, 448
- Nordhaus, J., & Blackman, E. G. 2006, *MNRAS*, 370, 2004
- Paltani, S., Türler, M. 2005, *A&A*, 435, 811
- Pelletier, G., & Pudritz, R. E. 1992, *ApJ*, 394, 117
- Pringle, J. E. 1981, *ARA&A*, 19, 137
- Shakura, N. I., Sunyaev, R. A., 1973, *A&A*, 24, 337
- Terquem, C., Eisloffel, J., Papaloizou, J. C. B., & Nelson, R. P. 1999, *ApJL*, 512, L131
- Terquem, C.E.J.M.L.J, 2002, *EAS Publications Series*, 3, 203
- Young, A. J., Fabian, A. C., Ross, R. R., & Tanaka, Y. 2001, *MNRAS*, 325, 1045
- Yuan, F., Markoff, S., Falcke, H., 2002, *A&A*, 383, 854